

**PHENYL-SUBSTITUTED INDOLES AND INDAZOLES**5                   **Cross-Reference to Related Applications**

This application claims priority from U.S. Provisional Application Serial Number 60/194,071, filed on March 31, 2000, our Docket Number ORT-1158 and U.S. Provisional Application Serial Number 60/194,071, filed on February 28, 2001, our Docket Number ORT-1366.

10

**Field of the Invention**

The invention relates to pharmaceutically-active fused heterobicyclic compounds and methods of using them to treat or prevent disorders and conditions, such as those mediated by the histamine H<sub>3</sub> receptor.

15

**Background**

The histamine H<sub>3</sub> receptor is located as a presynaptic autoreceptor in the central nervous system and as a presynaptic heteroreceptor on serotonergic, noradrenergic, dopaminergic, and cholinergic neurons. The histamine H<sub>3</sub> receptor is also located peripherally in tissues such as vascular smooth muscle cells.

Proposed uses of histamine H<sub>3</sub> antagonists include the treatment or prevention of dementia, Alzheimer's disease (Panula et al. *Abstr. Society Neuroscience*, 1995, 21:1977), epilepsy (Yokoyama et al. *Eur. J. Pharmacol.*, 1993, 234:129), sleep/wake disorders (Lin et al., *Br. Res.*, 1990, 523, 325; Monti et al., *Eur. J. Pharmacol.*, 1991, 205, 283) including narcolepsy, insomnia, and jet lag, eating disorders (Machidori et al. *Brain Research*, 1992, 590:180), motion sickness, vertigo, attention deficit hyperactivity disorder, learning and memory disorders (Barnes et al. *Abstr. Society Neuroscience*, 1993, 19:1813), schizophrenia (Schlicker et al. *Naunyn Schmiedeberg's Arch. Pharmacol.*, 1996, 353:325), and sequelae associated with post-ischemic reperfusion and hypertension. (Imamura et al., *J. Pharmacol. Expt. Ther.*, 1994, 271, 1259). H<sub>3</sub> antagonists are also useful to treat or prevent neurogenic


inflammation such as migraine (McLeod *et al.*, *Abstr. Society Neuroscience*, 1996, 22, 2010), asthma (Ichinose *et al.*, *Eur. J. Pharmacol.*, 989, 174, 49), obesity, allergic rhinitis, substance abuse, bipolar disorders, manic disorders, and depression. Histamine H<sub>3</sub> antagonists alone or in combination with a histamine H<sub>1</sub> antagonist are believed to be useful in the treatment of upper airway allergic response or allergic rhinitis (US Patent Nos. 5217986, 5352707, and 5869479).

As noted, the prior art related to histamine H<sub>3</sub> ligands was comprehensively reviewed recently (*"The Histamine H<sub>3</sub> Receptor-A Target for New Drugs"*, Leurs, R., and Timmerman, H., (Editors), Elsevier, 1998). Within this reference the medicinal chemistry of histamine H<sub>3</sub> agonists and antagonists was reviewed (see Krause *et al.* and Phillips *et al.*, respectively). Thus the importance of an imidazole moiety containing only a single substitution in the 4 position was noted together with the deleterious effects of additional substitution on activity. Particularly methylation of the imidazole ring at any of the remaining unsubstituted positions was reported to strongly decrease activity.

More recently several publications have described histamine H<sub>3</sub> ligands that do not contain an imidazole moiety. Examples include Ganellin *et al Arch. Pharm. (Weinheim, Ger.)* 1998, 331, 395; Walczynski *et al Arch. Pharm. (Weinheim, Ger.)* 1999, 332, 389; Walczynski *et al Farmaco* 1999, 684; Linney *et al J. Med. Chem.* 2000, 2362; U.S. Patent 5,352,707; PCT Application WO99/42458, published August 26, 1999; and European Patent Application 0978512, published on February 9, 2000.

### Summary of the Invention

The invention features phenyl-substituted indole and indazole compounds, methods of making them, and methods of using them. One aspect of the invention provides compounds of the following formula (I)(B):



5

wherein  $X_1$  is  $CR_1$ , wherein  $R_1$  is H, halo, cyano, amino, or nitro;  
and  $X_2$  is  $NR_3$ ;

$R_3$  is H,  $-SO_2$  ( $C_{1-6}$  alkyl),  $-SO_2$  phenyl,  $(C=O)(C_{1-6}$  alkyl), or  $-W'Z'$ ;

$W'$  is a covalent bond,  $(C=O)$ ,  $SO_2$ , or  $C_{1-6}$  alkyl;

10

$Z'$  is  $C_{1-6}$  alkyl,  $C_{1-6}$  alkoxy,  $C_{3-8}$  cycloalkyl, phenyl, or  $C_{2-6}$   
heterocyclic radical, optionally including in the ring up to 3 additional  
heteroatoms or moieties independently selected from O, N, NH, S, SO,  
and  $SO_2$ ; or  $Z'$  is  $NR_{13}R_{14}$  where each of  $R_{13}$  and  $R_{14}$  is independently  
selected from  $C_{1-6}$  alkyl,  $C_{2-6}$  alkenyl, phenyl, benzyl,  $C_{3-8}$  cycloalkyl,  
and  $C_{2-5}$  heterocyclic radical;

15

each of  $R_5$ ,  $R_6$ ,  $R_7$  and  $R_8$  is independently H,  $C_{1-6}$  alkyl,  $C_{1-6}$   
alkoxy, halo, nitro, or amino;

one of  $R_a$ ,  $R_b$ ,  $R_c$ ,  $R_d$ , and  $R_e$  is WZ and the others are  
independently selected from H,  $C_{1-6}$  alkyl,  $C_{1-6}$  alkoxy, halo, nitro, and  
amino;

20

W is  $-O-$ ,  $R_9$ ,  $O-R_9$ ,  $NR_{10}$ ,  $-(CO)(O)R_9$ ,  $-O(CO)R_9$ ,  
 $-(CO)NR_{10}$ , or  $-N(R_{10})-CO-R_9$ , wherein  $R_9$  is  $C_{1-6}$  alkylene,  $C_{2-6}$   
alkynylene,  $C_{2-6}$  alkenylene, phenylene, or  $C_{2-5}$  heterocyclic bivalent  
radical, and  $R_{10}$  is H,  $C_{1-6}$  alkyl,  $C_{2-6}$  alkynyl,  $C_{2-6}$  alkenyl, phenyl, or  $C_{2-5}$   
heterocyclic radical;

25

Z is  $C_{2-8}$  heterocyclic radical with at least one basic nitrogen atom  
in the ring, optionally including in the ring up to 3 additional heteroatoms  
or moieties independently selected from O,  $C=O$ , N, NH, NG, S, SO,  
and  $SO_2$ , wherein G is  $R_{15}$ ,  $COR_{15}$ ,  $COOR_{15}$ ,  $SO_2R_{15}$ ,  $SO_2N$ ,  $CSR_{15}$ ; or Z  
is  $NR_{11}R_{12}$  where each of  $R_{11}$  and  $R_{12}$  is independently selected from H,  
 $C_{1-6}$  alkyl, phenyl, benzyl,  $C_{3-8}$  cycloalkyl, and  $C_{2-5}$  heterocyclic radical;  
or  $NR_{11}R_{12}$  taken together is a  $C_{6-8}$  cycloalkylimino radical; and  $R_{15}$  is  $C_{1-6}$   
alkyl,  $C_{2-6}$  alkynyl,  $C_{2-6}$  alkenyl,  $C_{3-7}$  cycloalkyl, and  $C_{4-7}$  cycloalkenyl;

30

each of the above hydrocarbyl or heterocyclic groups being optionally substituted with between 1 and 3 substituents selected from C<sub>1-3</sub> alkyl, C<sub>1-3</sub> alkoxy, halo, hydroxy, phenyl, and phenyl(C<sub>1-3</sub> alkyl); and wherein each of the above heterocyclic groups may be attached to the rest of the molecule by a carbon atom or a heteroatom;  
 or a pharmaceutically acceptable salt, amide, ester, or hydrate thereof.

According to another aspect of the invention, the disclosed compounds and certain other compounds, are useful for the treatment and/or prevention of diseases and conditions mediated by the histamine 3 (H<sub>3</sub>) receptor.

A third aspect of the invention features methods of making the disclosed compounds.

Additional features of the invention are disclosed in the following description and examples, and in the appended claims.

### Detailed Description

The invention features pharmaceutically active phenyl-substituted  
5 indoles and indazoles and methods of making and using them. The  
description is organized as follows:

- A. Terms
- B. Compounds
- C. Synthetic Methods
- 10 D. Uses (including dosages, formulations, and related compounds)
- E. Synthetic Examples
- F. Biological Examples
- G. Other Embodiments
- H. Claims

15

#### A. Terms

The following terms are defined below and by their usage throughout  
this disclosure.

15 "Alkyl" includes straight chain and branched hydrocarbons with at least  
one hydrogen removed to form a radical group. Alkyl groups include methyl,  
ethyl, propyl, isopropyl, butyl, isobutyl, t-butyl, 1-methylpropyl, pentyl,  
isopentyl, sec-pentyl, hexyl, heptyl, octyl, and so on. Alkyl does not include  
cycloalkyl.

20 "Alkenyl" includes straight chain and branched hydrocarbon radicals as  
above with at least one carbon-carbon double bond ( $sp^2$ ). Alkenyls include  
ethenyl (or vinyl), prop-1-enyl, prop-2-enyl (or allyl), isopropenyl (or 1-  
methylvinyl), but-1-enyl, but-2-enyl, butadienyls, pentenyls, hexa-2,4-dienyl,  
and so on. Hydrocarbon radicals having a mixture of double bonds and  
triple bonds, such as 2-penten-4-ynyl, are grouped as alkynyls herein.

30 Alkenyl does not include cycloalkenyl.

"Alkynyl" include straight chain and branched hydrocarbon radicals as  
above with at least one carbon-carbon triple bond ( $sp$ ). Alkynyls include  
ethynyl, propynyls, butynyls, and pentynyls. Hydrocarbon radicals having a

mixture of double bonds and triple bonds, such as 2-penten-4-ynyl, are grouped as alkynyls herein. Alkynyl does not include cycloalkynyl.

"Alkoxy" includes a straight chain or branched alkyl group with a terminal oxygen linking the alkyl group to the rest of the molecule. Alkoxy includes methoxy, ethoxy, propoxy, isopropoxy, butoxy, t-butoxy, pentoxy and so on. "Aminoalkyl", "thioalkyl", and "sulfonylalkyl" are analogous to alkoxy, replacing the terminal oxygen atom of alkoxy with, respectively, NH (or NR), S, and SO<sub>2</sub>.

"Aryl" includes phenyl, naphthyl, biphenyl, and so on.

"Cycloalkyl" includes cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, and so on.

"Cycloalkenyl" includes cyclobutenyl, cyclobutadienyl, cyclopentenyl, cyclopentadienyl, cyclohexenyl, cyclohexadienyl, cyclohexatrienyl (phenyl), cycloheptenyl, and so on. "Cycloalkynyl" includes the analogous rings with one or more triple bonds.

"Heterocyclic radicals" include aromatic and nonaromatic rings having carbon atoms and at least one heteroatom (O, S, N) or heteroatom moiety (SO<sub>2</sub>, CO, CONH, COO) in the ring. Unless otherwise indicated, a heterocyclic radical may have a valence connecting it to the rest of the molecule through a carbon atom, such as 3-furyl or 2-imidazolyl, or through a heteroatom, such as N-piperidyl or 1-pyrazolyl. Examples of heterocyclic radicals include thiazoyl, furyl, pyran, isobenzofuranyl, pyrrolyl, imidazolyl, pyrazolyl, isothiazolyl, isoxazolyl, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, indolizyl, isoindolyl, indolyl, indazolyl, purinyl, quinolyl, furazanyl, pyrrolidinyl, pyrrolinyl, imidazolidinyl, imidazolinyl, pyrazolidinyl, pyrazolinyl, piperidyl, piperazinyl, indolinyl, and morpholinyl. For example, preferred heterocyclic radicals for Z include morpholinyl, piperazinyl, pyrazinyl, pyrrolidinyl, pyridyl, cyclohexylimino, cycloheptylimino, and more preferably, piperidyl.

"halo" includes fluoro, chloro, bromo, and iodo, and preferably fluoro or chloro.

"patient" or "subject" includes mammals such as humans and animals (dogs, cats, horses, rats, rabbits, mice, non-human primates) in need of

observation, experiment, treatment or prevention in connection with the relevant disease or condition. Preferably, the patient is a human.

"composition" includes a product comprising the specified ingredients in the specified amounts as well as any product which results directly or indirectly from combinations of the specified ingredients in the specified amounts.

Concerning the various radicals in this disclosure and in the claims, two general remarks are made. The first remark concerns valency. As with all hydrocarbon radicals, whether saturated, unsaturated or aromatic, and whether or not cyclic, straight chain, or branched, and also similarly with all heterocyclic radicals, each radical includes substituted radicals of that type and monovalent, bivalent, and multivalent radicals as indicated by the context of the claims. The context will indicate that the substituent is an alkylene or hydrocarbon radical with at least two hydrogen atoms removed (bivalent) or more hydrogen atoms removed (multivalent). An example of a bivalent radical linking two parts of the molecule is W in formula (I)(B) which links Z with the phenyl group (-Ph-WZ). Subject to the claims, W can be an alkyl (strictly, alkylene) group (-Ph-CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-Z), an aminoalkyl group (-Ph-NH-CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-Z), an alkoxy group (-Ph-O-CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-Z), an "oxa" (-Ph-O-Z), and so on.

Second, radicals or structure fragments as defined herein are understood to include substituted radicals or structure fragments. Using "alkyl" as an example, "alkyl" should be understood to include substituted alkyl having one or more substitutions, such as between 1 and 5, 1 and 3, or 2 and 4 substituents. The substituents may be the same (dihydroxy, dimethyl), similar (chlorofluoro), or different (chlorobenzyl- or aminomethyl-substituted). Examples of substituted alkyl include haloalkyl (such as fluoromethyl, chloromethyl, difluoromethyl, perchloromethyl, 2-bromoethyl, and 3-iodocyclopentyl), hydroxyalkyl, aminoalkyl, nitroalkyl, alkylalkyl, and so on.

Preferred substitutions for phenyl include methyl, methoxy, fluoromethyl, difluoromethyl, perfluoromethyl (trifluoromethyl), 1-fluoroethyl, 2-fluoroethyl,

ethoxy, fluoro, chloro, and bromo, and particularly methyl, fluoromethyl, perfluoro, methoxy, and fluoro.

- 5 Examples of other substituted radicals or fragments include 1-methyl-2-pyrrolidino, 4-(piperidyl)-piperidyl, [4-(N-benzyl)piperidyl]amino, 4-fluorobenzylamino, beta-hydroxyethoxy, beta-hydroxypropoxy, 2-oxo-pyrrolidino, 4-(1-methyl-4-piperidinyl), 4-(5-methyl-thiazoyl), 4-chlorobenzyl, 4-fluorobenzyl, 4-methylbenzyl, 4-methylpiperazinyl, piperazinyl, and 4-(1-isopropyl-4-piperidinyl).



## B. Compounds

One aspect of the invention features compounds of formula (I)B as described in the Summary section above. The invention encompasses the described compounds and pharmaceutically acceptable salts, esters, amides, and hydrates thereof.

Preferred compounds include those compounds of formula (I)B wherein: (a)  $R_3$  is H or  $C_{1-3}$  alkyl; (b)  $R_3$  is  $-(C=O)C_{1-6}$  alkyl; (c)  $R_3$  is  $-SO_2(C_{1-3}$  alkyl); (d)  $R_3$  is methylsulfonyl; (e)  $W'$  is a covalent bond; (f)  $W'$  is  $SO_2$  or  $(C=O)$ ; (g)  $R_c$  is WZ; (h)  $R_b$  or  $R_d$  is WZ; (i)  $W$  is ethoxy, propoxy, or butoxy; (j)  $W$  is  $-O-$ ; (k) one of  $R_b$ ,  $R_c$ , and  $R_e$  is WZ and the others are independently selected from H, methyl, ethyl, methoxy, ethoxy, amino, nitro, and halo; and  $R_a$  and  $R_d$  are each independently H or methyl; (l) at least two of the following apply:  $R_c$  is WZ;  $W$  is propoxy or ethoxy; and  $Z$  is N-piperidino, 2-(N-methyl)pyrrolidino, or N,N-dimethyl; or combinations thereof.

Additional preferred compounds include those wherein (m)  $Z$  is piperazino or N-methylpiperazino, and more preferably  $Z$  is pyrrolidino, N-methyl-pyrrolidino, pyridyl, thiazoyl, piperidino, or  $NR_{11}R_{12}$  where each of  $R_{11}$  and  $R_{12}$  is independently selected from H,  $C_{1-6}$  alkyl, phenyl, benzyl,  $C_{3-6}$  cycloalkyl, and  $C_{2-5}$  heterocyclic radical or taken together with the N form a  $C_8$  cycloalkylamino radical; or wherein (m) is combined with (a) through (l) above.

Further preferred compounds include those wherein (n) one of  $R_b$ ,  $R_c$ , and  $R_e$  is WZ and the others are independently selected from H, methyl, ethyl, methoxy, ethoxy, amino, and halo; and  $R_a$  and  $R_d$  are each independently H or methyl;  $W$  is  $-O-$  or  $C_{1-3}$  alkoxy;  $Z$  is piperazino or N-methylpiperazino, and more preferably pyrrolidino, N-methylpyrrolidino, pyridyl, thiazoyl, piperidino, or  $NR_{11}R_{12}$  where each of  $R_{11}$  and  $R_{12}$  is independently selected from H,  $C_{1-2}$  alkyl, phenyl, benzyl,  $C_{3-8}$  cycloalkyl, and  $C_{2-5}$  heterocyclic radical; each of  $R_6$  and  $R_7$  are each independently H, methyl, methoxy, or ethoxy; and each of  $R_5$  and  $R_8$  is H. Preferred compounds also include those wherein for example one or more of (a) through (n) is combined with (o)  $R_3$  is H or  $-SO_2(C_{1-6}$  alkyl); or (p)  $R_3$  is  $SO_2$ (phenyl) and  $(C=O)(C_{1-6}$  alkyl).

Examples of more preferred compounds include: (i) 2-[4-[2-[1-(methyl)-2-pyrrolidinyl]ethoxy]phenyl]-1H-indole; 2-[4-[2-[1-(methyl)-2-pyrrolidinyl]ethoxy]phenyl]-1-(methylsulfonyl)-1H-indole; 2-[4-[3-Piperidinopropoxy]phenyl]-1H-indole; 2-[4-[3-(methylsulfonyl)-1H-indole]; and 2-[3-[3-Piperidinopropoxy]phenyl]-1-(methylsulfonyl)-1H-indole; and (ii) 2-(4-(3-(4-methylpiperazino)propoxy)-phenyl)indole; and 1-(methylsulfonyl)-2-(4-(3-(4-methylpiperazino)-propoxy)phenyl)indole.

Other examples of compounds, and methods of making them, are provided in the next section.

### C. Synthetic Chemical Methods


The invention provides methods of making the disclosed compounds according to traditional organic synthetic methods as well as matrix or combinatorial synthetic methods. Schemes 1 through 9 describe suggested synthetic routes.

Using these Schemes, the guidelines below, and the examples in section E, a person of skill in the art may develop analogous or similar methods for a given compound.

One skilled in the art will recognize that synthesis of the compounds of the present invention may be effected by purchasing an intermediate or protected intermediate compounds described in any of the schemes disclosed herein. One skilled in the art will further recognize that during any of the processes for preparation of the compounds in the present invention, it may be necessary and/or desirable to protect sensitive or reactive groups on any of the molecules concerned. This may be achieved by means of conventional protecting groups, such as those described in "Protective Groups in Organic Synthesis", John Wiley & Sons, 1991. These protecting groups may be removed at a convenient stage using methods known from the art.

Examples of the described synthetic routes includes Synthetic Examples 1 through 17. Compounds analogous to the target compounds of these examples can be, and in many cases, have been, made according to similar routes. The disclosed compounds are useful in basic research and as pharmaceutical agents as described in the next section.


Scheme I



- 5 Generally, a compound of formula (V), a known compound or compound prepared by known methods, is reacted with a compound of formula (II), a known compound or compound prepared by known methods, in the presence of a palladium catalyst such as dichlorobis(triphenylphosphine) palladium, and

CuI, or the like, in the presence of a base such as triethylamine, or the like, in a solvent such as DMF, THF, DMA, and the like, to yield the corresponding compound of formula (IV). Compound (IV) is then further reacted, as outlined in Schemes 5-7 below, to form the compound of formula (I). Alternatively a compound of formula (III) can be reacted with a compound of formula (II) using the above described, or similar methods to form a compound of formula (I) directly. In addition compounds of formula (I), in which  $X_2$  is NH can be obtained by reacting a compound of formula (VI) with an aromatic hydrazine of formula (VII) in the presence of a strong acid such as PPA.

Compounds of formula (II) wherein  $R_3$  is chosen from either  $-\text{SO}_2(\text{C}_{1-6} \text{ alkyl})$ ,  $-\text{SO}_2$  phenyl,  $(\text{C}=\text{O})(\text{C}_{1-6} \text{ alkyl})$ , and  $R_5-R_8$  are selected from H,  $\text{C}_{1-6}$  alkyl,  $\text{C}_{1-6}$  alkoxy, halo, nitro, may be prepared according to the process outlined in Scheme 2.




A compound of formula (VIII), wherein  $R_5-R_8$  are selected from H,  $\text{C}_{1-6}$  alkyl,  $\text{C}_{1-6}$  alkoxy, halo, nitro, a known compound or compound prepared by known methods, is treated with an iodinating agent such as N-iodo succinamide, ICl, or  $\text{I}_2$  in a solvent such as acetic acid, acetonitrile, or the like, to yield the corresponding compound of formula (IX). The compound (IX), a known compound or compound prepared by known methods, is reacted with a compound of formula (XI), in which  $R_3$  is chosen from either  $-\text{SO}_2(\text{C}_{1-6} \text{ alkyl})$ ,  $-\text{SO}_2$ phenyl,  $(\text{C}=\text{O})(\text{C}_{1-6} \text{ alkyl})$ ,  $(\text{C}=\text{O})(\text{C}_{1-6} \text{ alkoxy})$ ,  $(\text{C}=\text{O})$ phenyl, and  $X_3$  is selected from Br, Cl, F, or a conventional activating anhydride, or ester, in the presence of a base such as pyridine, N,N-dimethyl aminopyridine, triethylamine, or sodium hydroxide in an organic solvent such as DCM, THF, or DMF to yield the corresponding compound of formula (II).

Compounds of formula (V) wherein W is OH, NH<sub>2</sub>, CO<sub>2</sub>H, and Ra-Re are selected from H, C<sub>1-6</sub> alkyl, C<sub>1-6</sub> alkoxy, halo, or nitro may be prepared according to the processes outlined in Scheme 3.

5


Scheme 3



A compound of formula (X) wherein W is OH, NH<sub>2</sub>, CO<sub>2</sub>H, and Ra-Re are selected from H, C<sub>1-6</sub> alkyl, C<sub>1-6</sub> alkoxy, halo, or nitro, is reacted with a  
 10 diazophosphonate in the presence of a base such as, K<sub>2</sub>CO<sub>3</sub>, KOH, or DBU, in a solvent such as MeOH, EtOH, or DMF, to yield compounds of formula (V). Alternately compounds of formula (V) can also be prepared by treating a compound of formula (X) with trimethylsilyldiazomethane, in the presence of a  
 15 strong base such as LDA or LHMDs, in a solvent such as, THF, Ether, or MTBE, to yield compounds of formula (V). In addition compounds of formula (V) may also be obtained using methods known to one skilled in the art as outlined in R. C. Larock "Comprehensive Organic Transformations", VCH Publishers, 1989, p. 295-296.

Specifically compounds of formula (IV) wherein W is -OH, -NH<sub>2</sub>, -  
 20 C(O)OH may be prepared according to the processes outlined in Scheme 4.

Scheme 4




25 Specifically, Compounds of formula (II), as defined in Scheme 2, are combined with compounds of formula (V), as defined in Scheme 3, in the

presence of a palladium catalyst such as,  $\text{Pd}(\text{PPh}_3)_2\text{Cl}_2$ , or  $\text{Pd}(\text{OAc})_2$ , and a copper source such as  $\text{CuI}$ ,  $\text{CuOAc}$ , or  $\text{CuBr}$ , and a base such as triethylamine or pyridine, in a solvent such as THF or DMF, to provide the corresponding compounds of formula (IV).

- 5           Compounds of formula (I) in which  $n$  is a whole number between 0 and 4, and  $Z$  is as described in claim (1), and  $R_5$ - $R_8$  and  $R_a$ - $R_e$  are as described in Scheme 4, can be obtained by the procedures described in Scheme 5-7.

Scheme 5




- Specifically a compound of formula (IVa) wherein  $R_3$ ,  $R_a$ - $R_e$  and  $R_5$ - $R_8$  are as described in Scheme 4, is reacted with an alcohol of formula (XII), wherein  $Z$  is as described in claim (1), and  $n$  is a whole number between 0 and 4, in the presence of a phosphine such as triphenyl phosphine, polymer supported triphenylphosphine, or tributylphosphine, and an azodicarboxylate such as diisopropylazodicarboxylate, 1,1'-(azodicarbonyl)dipiperidine, or other Mitsunobu conditions, in a solvent such as DCM or THF, to afford the corresponding compounds of formula (I) in which  $X_4$  is a covalent bond, and  $n$  is a whole number between 0 and 4.

- Alternatively compounds of formula (IV) as described above, can be reacted with carboxylic acids of formula (XIII), in which  $Z$  is defined as above, and  $n$  is a whole number between 0 and 3, in the presence of an activating agent such as carbonyldiimidazole or thionyl chloride, with a base such as  $N$ -methyl morpholine, triethylamine, or  $N,N$ -dimethyl-4-aminopyridine to yield the corresponding compound of formula (I), in which  $X_4$  is defined as a carbonyl group.

Alternatively compounds of formula (Ib) can be obtained by the methods described in Scheme 6.


Scheme 6



Specifically, a compound of formula (IVb) in which R<sub>3</sub>, R<sub>5</sub>-R<sub>8</sub> and R<sub>a</sub>-R<sub>e</sub> is as defined in scheme 4, is reacted with an aldehyde of formula (XIV) in which n is a whole number between 0 and 3, and Z is as described in claim (1), in the presence of a reducing agent such as NaBH<sub>3</sub>(CN) or NaBH(OAc)<sub>3</sub>, in a solvent such as MeOH or THF, to afford the corresponding compound of formula (I).

Alternatively, compounds of formula (Ic) can be obtained using the methods outlined in Scheme 7.

Scheme 7




Specifically, a compound of formula (IVc) in which R<sub>3</sub>, R<sub>5</sub>-R<sub>8</sub>, and R<sub>a</sub>-R<sub>e</sub> is defined as in Scheme 4, is reacted with an alcohol of formula (XII), or an amine of formula (XV), in which Z is as defined in Claim (1), and n is a whole number between 0 and 4, in the presence of an activating agent such as carbonyldiimidazole or thionyl chloride, with a base such as N-methyl morpholine, triethylamine, or N,N-dimethyl-4-aminopyridine to yield the corresponding compound of formula (I), in which X<sub>5</sub> is defined as O or NH.



In addition, compounds of formula (I)B can be converted to other compounds of formula (I)B as defined in Scheme 8 below.

Scheme 8



Specifically, a compound of formula (Id) in which  $R_3$ ,  $R_5\text{-}R_8$ , and  $R_a\text{-}R_e$  are as described in Scheme 4, is treated with; a nitrating agent such as  $\text{HNO}_3$  or an electrophilic halogenating agent such as  $\text{Br}_2$  or  $\text{NIS}$ , using solvents and conditions known to one skilled in the art, to yield the corresponding compound of formula (If) in which  $R_1$  is defined as  $\text{NO}_2$ ,  $\text{Br}$ ,  $\text{Cl}$ , or  $\text{I}$ . Additionally a compound of formula (If) in which  $R_1$  is  $\text{NO}_2$  can be further elaborated through reduction with an appropriate reducing agent such as  $\text{SnCl}_2$  or iron metal, to yield the corresponding compound of formula (Ie), in which  $R_1$  is  $\text{NH}_2$ .


Additionally, compounds of formula (Ie) in which  $R_3$  is defined as in Scheme 2, and  $R_5\text{-}R_8$ , and  $R_a\text{-}R_e$  are as described in Scheme 4, can be treated with a strong base such as  $\text{KOH}$ ,  $\text{K}_2\text{CO}_3$ , or the like, in a solvent such as THF, MeOH, or the like, to yield the corresponding compounds of formula (If).

In addition, compounds of formula (If) can be converted to compounds of the corresponding formula (Ie) by treatment with a strong base such as  $n\text{-BuLi}$ ,  $\text{NaH}$ , or the like, and an alkylating or acylating agent of formula (XI), wherein  $R_3$  and  $X_3$  are as defined in Scheme 2.

The synthesis of compounds of formula (III) in which  $R_a\text{-}R_e$  are as defined in Scheme 3,  $R_{11}$ , and  $R_{12}$  are as defined claim (1), and  $n$  is an integer

[illegible]

Scheme 9




Specifically, a compound of formula (XVI) in which  $R_{13}$  is H, or  $C_{1-6}$  alkyl, and  $R_a-R_e$  is as previously described, is treated with a base such as NaH or  $K_2CO_3$ , and a compound of formula (XVIII), in which  $X_6$  is selected from Cl, Br, I,  $-OSO_2CH_3$ ,  $-OTs$ , or  $OTf$ , and  $X_7$  is selected from the same definition as  $X_6$  but less reactive than the element chosen for  $X_6$ , and  $n$  is an integer from 2 to 5, in a solvent such as THF, DMF or DMSO, to yield the corresponding compound of formula (XVII). The compound of formula (XVII) is then treated with a compound of formula (XIX), wherein  $R_{11}$  and  $R_{12}$  are as defined in Claim (1), in a solvent such as DMF or DCM, to afford the corresponding compound of formula (VI).

Compounds of formula (III) are prepared by treatment of corresponding compounds of formula (IV), in which  $R_{13}$  is defined as H, with a base such as LDA or  $LiHMDS$ , and  $TMSCHN_2$ , in a solvent such as THF, diethylether, or the like. Alternately, compounds of formula (III) can also be prepared by treating corresponding compounds of formula (IV), with a base such as  $K_2CO_3$  or  $KOH$ , and a phosphonate such as  $(CH_3O)_2P(O)C(N_2)C(O)CH_3$ , in a solvent such as MeOH.

Compounds of formula (Ig) can be obtained using the methods described in Scheme 10.


Scheme 10



5 Specifically a compound of formula (VI) as defined in scheme 9 is treated with an aryl hydrazine of formula (VII), wherein R<sub>5</sub>-R<sub>8</sub> is as defined as in claim (1), in polyphosphoric acid, to yield the corresponding compound of formula (Ig).

10 Additionally compounds of formula (I) can be formed using the procedures outlined in Scheme 11.

Scheme 11



15 Specifically a compound of formula (II), as defined in Scheme 2 is combined with a compound of formula (III) as defined in Scheme 9, in the presence of a palladium catalyst such as Pd(PPh<sub>3</sub>)<sub>2</sub>Cl<sub>2</sub> or Pd(OAc)<sub>2</sub>, and a copper source such as CuI, CuOAc or CuBr, and a base such as triethylamine or pyridine, in a solvent such as THF or DMF, to provide the corresponding compounds of formula (Ih).

20

#### D. Uses

25 According to the invention, the disclosed compounds and compositions are useful for the amelioration of symptoms associated with, the treatment of, and/or the prevention of, the following conditions and diseases, or symptoms associated with them: dementia, Alzheimer's disease, narcolepsy, eating disorders, motion sickness, vertigo, attention deficit hyperactivity disorder,

learning and memory disorders, schizophrenia, mild cognitive impairment upper airway allergic response (allergic rhinitis), insomnia, jet lag, obesity, asthma, neurogenic inflammation, substance abuse, bipolar disorders, manic disorders, and depression. The invention also features pharmaceutical compositions, which include, without limitation, one or more of the disclosed compounds, and a pharmaceutically acceptable carrier or excipient.

### 1. Dosages

Those skilled in the art will be able to determine, according to known methods, the appropriate dosage for a patient, taking into account factors such as age, weight, general health, the type of symptoms requiring treatment, and the use of other medications. An effective amount means that amount of pharmaceutical reagent (such as a prodrug, metabolic precursor, or active compound) that elicits the biological or medical response desired. In general, a therapeutically effective amount will be between 0.01 and 1000 mg/kg per day, preferably between 0.01 and 250 mg/kg body weight, and daily dosages will be between 0.50 and 5000 mg for an adult subject of normal weight. Capsules, tablets or other formulations (such as liquids and film-coated tablets) may be of between 0.20 and 100 mg, such as 0.20, 0.50, 1.0, 2.0, 3.0, and 10 mg and can be administered according to the disclosed methods.

### 2. Formulations

Dosage unit forms include tablets, capsules, pills, powders, granules, aqueous and nonaqueous oral solutions and suspensions, and parenteral solutions packaged in containers adapted for subdivision into individual doses. Dosage unit forms can also be adapted for various methods of administration, including controlled release formulations, such as subcutaneous implants. Administration methods include oral, rectal, parenteral (intravenous, intramuscular, subcutaneous), intracisternal, intravaginal, intraperitoneal, intravesical, local (drops, powders, ointments, gels or cream), and by inhalation (a buccal or nasal spray) as appropriate depending on the overall health and condition of the patient as determined by a physician or veterinary doctor.

Parenteral formulations include pharmaceutically acceptable aqueous or nonaqueous solutions, dispersion, suspensions, emulsions, and sterile powders for the preparation thereof. Examples of carriers include water,

ethanol, polyols (propylene glycol, polyethylene glycol), vegetable oils, and injectable organic esters such as ethyl oleate. Fluidity can be maintained by the use of a coating such as lecithin, a surfactant, or maintaining appropriate particle size. Carriers for solid dosage forms include (a) fillers or extenders, (b) binders, (c) humectants, (d) disintegrating agents, (e) solution retarders, (f) absorption accelerators, (g) adsorbants, (h) lubricants, (i) buffering agents, and (j) propellants.

Compositions may also contain adjuvants such as preserving, wetting, emulsifying, and dispensing agents; antimicrobial agents such as parabens, chlorobutanol, phenol, and sorbic acid; isotonic agents such as a sugar or sodium chloride; absorption-prolonging agents such as aluminum monostearate and gelatin; and absorption-enhancing agents.

### 3. Combination Therapy

The present invention also provides compositions and methods useful for the treatment of disorders or conditions modulated, preferably antagonized, by the histamine  $H_3$  receptor in combination with compounds that modulate other receptors including, but not limited to, histamine  $H_1$  and histamine  $H_2$  receptors. The present invention includes compounds and compositions useful in methods of combination therapy for the treatment of diseases or conditions modulated by the histamine  $H_3$  receptor in combination with compounds that are selective serotonin re-uptake inhibitors (SSRIs), such as PROZAC™, or are selective norepinephrine uptake inhibitors. Such combination methods include (a) administering the two or more pharmaceutical agents separately formulated and at separate times, and (b) administering the two or more agents simultaneously in a single formulation or in separate formulations administered more or less at the same time. For example, one aspect is a method of treatment comprising administering at least one histamine  $H_3$  receptor modulating compound disclosed herein and administering at least one compound selected from a histamine  $H_1$  receptor modulating compound, a histamine  $H_2$  receptor modulating compound, a selective serotonin reuptake inhibitor (such as PROZAC™), or a selective norepinephrine uptake inhibiting compound.

#### 4. Related Compounds

The invention provides the disclosed compounds and closely related, pharmaceutically acceptable forms of the disclosed compounds, such as salts, esters, amides, acids, hydrates or solvated forms thereof; masked or protected forms; and racemic mixtures, or enantiomerically or optically pure forms.

Pharmaceutically acceptable salts, esters, and amides include carboxylate salts (e.g., C<sub>1-8</sub> alkyl, cycloalkyl, aryl, heteroaryl, or non-aromatic heterocyclic) amino acid addition salts, esters, and amides which are within a reasonable benefit/risk ratio, pharmacologically effective and suitable for contact with the tissues of patients without undue toxicity, irritation, or allergic response. Representative salts include hydrobromide, hydrochloride, sulfate, bisulfate, nitrate, acetate, oxalate, valerate, oleate, palmitate, stearate, laurate, borate, benzoate, lactate, phosphate, tosylate, citrate, maleate, fumarate, succinate, tartrate, naphthylate, mesylate, glucoheptonate, lactobionate, and laurylsulfonate. These may include alkali metal and alkali earth cations such as sodium, potassium, calcium, and magnesium, as well as non-toxic ammonium, quaternary ammonium, and amine cations such as tetramethyl ammonium, methylamine, trimethylamine, and ethylamine. See example, S.M. Berge, et al., "Pharmaceutical Salts," J. Pharm. Sci., 1977, 66:1-19 which is incorporated herein by reference. Representative pharmaceutically acceptable amides of the invention include those derived from ammonia, primary C<sub>1-6</sub> alkyl amines and secondary di (C<sub>1-6</sub> alkyl) amines. Secondary amines include 5- or 6-membered heterocyclic or heteroaromatic ring moieties containing at least one nitrogen atom and optionally between 1 and 2 additional heteroatoms. Preferred amides are derived from ammonia, C<sub>1-3</sub> alkyl primary amines, and di (C<sub>1-2</sub> alkyl)amines. Representative pharmaceutically acceptable esters of the invention include C<sub>1-7</sub> alkyl, C<sub>5-7</sub> cycloalkyl, phenyl, and phenyl(C<sub>1-6</sub>)alkyl esters. Preferred esters include methyl esters.

The invention also includes disclosed compounds having one or more functional groups (e.g., hydroxyl, amino, or carboxyl) masked by a protecting group. See, e.g., Greene and Wuts, Protective Groups in Organic Synthesis, 3<sup>rd</sup> ed., (1999) John Wiley & Sons, NY. Some of these masked or protected

	1970	1971	1972	1973	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	2026	2027	2028	2029	2030	2031	2032	2033	2034	2035	2036	2037	2038	2039	2040	2041	2042	2043	2044	2045	2046	2047	2048	2049	2050	2051	2052	2053	2054	2055	2056	2057	2058	2059	2060	2061	2062	2063	2064	2065	2066	2067	2068	2069	2070	2071	2072	2073	2074	2075	2076	2077	2078	2079	2080	2081	2082	2083	2084	2085	2086	2087	2088	2089	2090	2091	2092	2093	2094	2095	2096	2097	2098	2099	2100	2101	2102	2103	2104	2105	2106	2107	2108	2109	2110	2111	2112	2113	2114	2115	2116	2117	2118	2119	2120	2121	2122	2123	2124	2125	2126	2127	2128	2129	2130	2131	2132	2133	2134	2135	2136	2137	2138	2139	2140	2141	2142	2143	2144	2145	2146	2147	2148	2149	2150	2151	2152	2153	2154	2155	2156	2157	2158	2159	2160	2161	2162	2163	2164	2165	2166	2167	2168	2169	2170	2171	2172	2173	2174	2175	2176	2177	2178	2179	2180	2181	2182	2183	2184	2185	2186	2187	2188	2189	2190	2191	2192	2193	2194	2195	2196	2197	2198	2199	2200	2201	2202	2203	2204	2205	2206	2207	2208	2209	2210	2211	2212	2213	2214	2215	2216	2217	2218	2219	2220	2221	2222	2223	2224	2225	2226	2227	2228	2229	2230	2231	2232	2233	2234	2235	2236	2237	2238	2239	2240	2241	2242	2243	2244	2245	2246	2247	2248	2249	2250	2251	2252	2253	2254	2255	2256	2257	2258	2259	2260	2261	2262	2263	2264	2265	2266	2267	2268	2269	2270	2271	2272	2273	2274	2275	2276	2277	2278	2279	2280	2281	2282	2283	2284	2285	2286	2287	2288	2289	2290	2291	2292	2293	2294	2295	2296	2297	2298	2299	2300	2301	2302	2303	2304	2305	2306	2307	2308	2309	2310	2311	2312	2313	2314	2315	2316	2317	2318	2319	2320	2321	2322	2323	2324	2325	2326	2327	2328	2329	2330	2331	2332	2333	2334	2335	2336	2337	2338	2339	2340	2341	2342	2343	2344	2345	2346	2347	2348	2349	2350	2351	2352	2353	2354	2355	2356	2357	2358	2359	2360	2361	2362	2363	2364	2365	2366	2367	2368	2369	2370	2371	2372	2373	2374	2375	2376	2377	2378	2379	2380	2381	2382	2383	2384	2385	2386	2387	2388	2389	2390	2391	2392	2393	2394	2395	2396	2397	2398	2399	2400	2401	2402	2403	2404	2405	2406	2407	2408	2409	2410	2411	2412	2413	2414	2415	2416	2417	2418	2419	2420	2421	2422	2
--	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	---



HYDROXYL PROTECTING GROUPS

- Protection for the hydroxyl group includes methyl ethers, substituted methyl ethers, substituted ethyl ethers, substitute benzyl ethers, and silyl ethers.

Substituted Methyl Ethers

- Examples of substituted methyl ethers include methoxymethyl, methylthiomethyl, *t*-butylthiomethyl, (phenyldimethylsilyl)methoxymethyl, benzyloxymethyl, *p*-methoxybenzyloxymethyl, (4-methoxyphenoxy)methyl, guaiacolmethyl, *t*-butoxymethyl, 4-pentenylloxymethyl, siloxymethyl, 2-methoxyethoxymethyl, 2,2,2-trichloroethoxymethyl, bis(2-chloroethoxy)methyl, 2-(trimethylsilyl)ethoxymethyl, tetrahydropyranyl, 3-bromotetrahydropyranyl, tetrahydrothiopyranyl, 1-methoxycyclohexyl, 4-methoxytetrahydropyranyl, 4-methoxytetrahydrothiopyranyl, 4-methoxytetrahydrothiopyranyl S,S-dioxido, 1-[(2-chloro-4-methyl)phenyl]-4-methoxypiperidin-4-yl, 1,4-dioxan-2-yl, tetrahydrofuranyl, tetrahydrothiofuranyl and 2,3,3a,4,5,6,7,7a-octahydro-7,8,8-trimethyl-4,7-methanobenzofuran-2-yl.

Substituted Ethyl Ethers

- Examples of substituted ethyl ethers include 1-ethoxyethyl, 1-(2-chloroethoxy)ethyl, 1-methyl-1-methoxyethyl, 1-methyl-1-benzyloxyethyl, 1-methyl-1-benzyloxy-2-fluoroethyl, 2,2,2-trichloroethyl, 2-trimethylsilylethyl, 2-(phenylselenyl)ethyl, *t*-butyl, allyl, *p*-chlorophenyl, *p*-methoxyphenyl, 2,4-dinitrophenyl, and benzyl.

Substituted Benzyl Ethers

- Examples of substituted benzyl ethers include *p*-methoxybenzyl, 3,4-dimethoxybenzyl, *o*-nitrobenzyl, *p*-nitrobenzyl, *p*-halobenzyl, 2,6-dichlorobenzyl, *p*-cyanobenzyl, *p*-phenylbenzyl, 2- and 4-picolyl, 3-methyl-2-picolyl N-oxido, diphenylmethyl, *p*, *p'*-dinitrobenzhydryl, 5-dibenzosuberyl, triphenylmethyl,  $\alpha$ -naphthyldiphenylmethyl, *p*-methoxyphenyldiphenylmethyl, di(*p*-methoxyphenyl)phenylmethyl, tri(*p*-methoxyphenyl)methyl, 4-(4'-

- bromophenacyloxy)phenyldiphenylmethyl, 4,4',4''-tris(4,5-dichlorophthalimidophenyl)methyl, 4,4',4''-tris(levulinoyloxyphenyl)methyl, 4,4',4''-tris(benzoyloxyphenyl)methyl, 3-(1-midazol-1-ylmethyl)bis(4',4''-dimethoxyphenyl)methyl, 1,1-bis(4-methoxyphenyl)-1'-pyrenylmethyl, 9-anthryl, 9-(9-phenyl)xanthenyl, 9-(9-phenyl-10-oxo)anthryl, 1,3-benzodithiolan-2-yl, and 5 benzisothiazolyl S,S-dioxido.

### Silyl Ethers

- Examples of silyl ethers include trimethylsilyl, triethylsilyl, triisopropylsilyl, 10 dimethylisopropylsilyl, diethylisopropylsilyl, dimethylhexylsilyl, *t*-butyldimethylsilyl, *t*-butyldiphenylsilyl, tribenzylsilyl, tri-*p*-xylylsilyl, triphenylsilyl, diphenylmethylsilyl, and *t*-butylmethoxyphenylsilyl.

### Esters

- 15 In addition to ethers, a hydroxyl group may be protected as an ester. Examples of esters include formate, benzoylformate, acetate, chloroacetate, dichloroacetate, trichloroacetate, trifluoroacetate, methoxyacetate, triphenylmethoxyacetate, phenoxyacetate, *p*-chlorophenoxyacetate, *p*-P-phenylacetate, 3-phenylpropionate, 4-oxopentanoate(levulinate), 4,4- 20 (ethylenedithio)pentanoate, pivaloate, adamantate, crotonate, 4-methoxycrotonate, benzoate, *p*-phenylbenzoate, 2,4,6-trimethylbenzoate(mesitoate)

### Carbonates

- 25 Examples of carbonate protecting groups include methyl, 9-fluorenylmethyl, ethyl, 2,2,2-trichloroethyl, 2-(trimethylsilyl)ethyl, 2-(phenylsulfonyl)ethyl, 2-(triphenylphosphonio)ethyl, isobutyl, vinyl, allyl, *p*-nitrophenyl, benzyl, *p*-methoxybenzyl, 3,4-dimethoxybenzyl, *o*-nitrobenzyl, *p*-nitrobenzyl, S-benzyl thiocarbonate, 4-ethoxy-1-naphthyl, and methyl 30 dithiocarbonate.

### Assisted Cleavage

Examples of assisted cleavage include 2-iodobenzoate, 4-azidobutyrate, 4-nitro-4-methylpentanoate, *o*-(dibromomethyl)benzoate, 2-formylbenzenesulfonate, 2-(methylthiomethoxy)ethyl carbonate, 4-(methylthiomethoxy)butyrate, and 2-(methylthiomethoxymethyl)benzoate.

### Miscellaneous Esters

Examples of miscellaneous esters include 2,6-dichloro-4-methylphenoxyacetate, 2,6-dichloro-4-(1,1,3,3-tetramethylbutyl)phenoxyacetate, 2,4-bis(1,1-dimethylpropyl)phenoxyacetate, chlorodiphenylacetate, isobutyrate, monosuccinoate, (E)-2-methyl-2-butenolate(tigloate), *o*-(methoxycarbonyl)benzoate, *p*-P-benzoate,  $\alpha$ -naphthoate, nitrate, alkyl N,N,N',N'-tetramethylphosphorodiamidate, N-phenylcarbamate, borate, dimethylphosphinothioyl, and 2,4-dinitrophenylsulfenate

### Sulfonates

Examples of sulfonates include sulfate, methanesulfonate(mesylate), benzyisulfonate, and tosylate.

## AMINO PROTECTING GROUPS

Protection for the amino group includes carbamates, amides, and special -NH protective groups.

Examples of carbamates include methyl and ethyl carbamates, substituted ethyl carbamates, assisted cleavage carbamates, photolytic cleavage carbamates, urea-type derivatives, and miscellaneous carbamates.

### Carbamates

Examples of methyl and ethyl carbamates include methyl and ethyl, 9-fluorenylmethyl, 9-(2-sulfo)fluorenylmethyl, 9-(2,7-dibromo)fluorenylmethyl, 2,7-

di-*t*-butyl-[9-(10,10-dioxo-10,10,10,10-tetrahydrothioxanthyl)]methyl, and 4-methoxyphenacyl.

#### Substituted Ethyl

5        Examples of substituted ethyl carbamates include 2,2,2-trichloroethyl, 2-trimethylsilylethyl, 2-phenylethyl, 1-(1-adamantyl)-1-methylethyl, 1,1-dimethyl-2-haloethyl, 1,1-dimethyl-2,2-dibromoethyl, 1,1-dimethyl-2,2,2-trichloroethyl, 1-methyl-1-(4-biphenyl)ethyl, 1-(3,5-di-*t*-butylphenyl)-1-methylethyl, 2-(2'- and 4'-pyridyl)ethyl, 2-(*N,N*-dicyclohexylcarboxamido)ethyl, *t*-butyl, 1-adamantyl, 10 vinyl, allyl, 1-isopropylallyl, cinnamyl, 4-nitrocinnamyl, 8-quinolyl, *N*-hydroxypiperidinyl, alkylidithio, benzyl, *p*-methoxybenzyl, *p*-nitrobenzyl, *p*-bromobenzyl, *p*-chlorobenzyl, 2,4-dichlorobenzyl, 4-methylsulfinylbenzyl, 9-anthrylmethyl and diphenylmethyl.

#### Assisted Cleavage

15        Examples of assisted cleavage include 2-methylthioethyl, 2-methylsulfonylethyl, 2-(*p*-toluenesulfonyl)ethyl, [2-(1,3-dithianyl)]methyl, 4-methylthiophenyl, 2,4-dimethylthiophenyl, 2-phosphonioethyl, 2-triphenylphosphonioisopropyl, 1,1-dimethyl-2-cyanoethyl, *m*-chloro-*p*-acyloxybenzyl, *p*-(dihydroxyboryl)benzyl, 5-benzisoxazolylmethyl, and 2-20 (trifluoromethyl)-6-chromonylmethyl.

#### Photolytic Cleavage

25        Examples of photolytic cleavage include *m*-nitrophenyl, 3,5-dimethoxybenzyl, *o*-nitrobenzyl, 3,4-dimethoxy-6-nitrobenzyl, and phenyl(*o*-nitrophenyl)methyl.

#### Urea-Type Derivatives

30        Examples of urea-type derivatives include phenothiazinyl-(10)-carbonyl derivative, *N'*-*p*-toluenesulfonylaminocarbonyl, and *N'*-phenylaminothiocarbonyl.

#### Miscellaneous Carbamates

Examples of miscellaneous carbamates include *t*-amyl, *S*-benzyl thiocarbamate, *p*-cyanobenzyl, cyclobutyl, cyclohexyl, cyclopentyl, cyclopropylmethyl, *p*-decyloxybenzyl, diisopropylmethyl, 2,2-dimethoxycarbonylvinyl, *o*-(*N,N*-dimethylcarboxamido)benzyl, 1,1-dimethyl-3-  
 5 (N,N-dimethylcarboxamido)propyl, 1,1-dimethylpropynyl, di(2-pyridyl)methyl, 2-furanylmethyl, 2-iodoethyl, isobornyl, isobutyl, isonicotinyl, *p*-(*p*'-methoxyphenylazo)benzyl, 1-methylcyclobutyl, 1-methylcyclohexyl, 1-methyl-1-cyclopropylmethyl, 1-methyl-1-(3,5-dimethoxyphenyl)ethyl, 1-methyl-1-(*p*-phenylazophenyl)ethyl, 1-methyl-1-phenylethyl, 1-methyl-1-(4-pyridyl)ethyl,  
 10 phenyl, *p*-(phenylazo)benzyl, 2,4,6-tri-*t*-butylphenyl, 4-(trimethylammonium)benzyl, and 2,4,6-trimethylbenzyl.

Examples of amides include:

#### 15 Amides

*N*-formyl, *N*-acetyl, *N*-chloroacetyl, *N*-trichloroacetyl, *N*-trifluoroacetyl, *N*-phenylacetyl, *N*-3-phenylpropionyl, *N*-picolinoyl, *N*-3-pyridylcarboxamide, *N*-benzoylphenylalanyl derivative, *N*-benzoyl, *N*-*p*-phenylbenzoyl.

#### 20 Assisted Cleavage

*N*-*o*-nitrophenylacetyl, *N*-*o*-nitrophenoxyacetyl, *N*-acetoacetyl, (*N*'-dithiobenzyloxycarbonylamino)acetyl, *N*-3-(*p*-hydroxyphenyl)propionyl, *N*-3-(*o*-nitrophenyl)propionyl, *N*-2-methyl-2-(*o*-nitrophenoxy)propionyl, *N*-2-methyl-2-(*o*-phenylazophenoxy)propionyl, *N*-4-chlorobutyryl, *N*-3-methyl-3-nitrobutyryl,  
 25 *N*-*o*-nitrocinnamoyl, *N*-acetylmethionine derivative, *N*-*o*-nitrobenzoyl, *N*-*o*-(benzoyloxymethyl)benzoyl, and 4,5-diphenyl-3-oxazolin-2-one.

#### Cyclic Imide Derivatives

*N*-phthalimide, *N*-dithiasuccinoyl, *N*-2,3-diphenylmaleoyl, *N*-2,5-dimethylpyrrolyl, *N*-1,1,4,4-tetramethyldisilylazacyclopentane adduct, 5-substituted 1,3-dimethyl-1,3,5-triazacyclohexan-2-one, 5-substituted 1,3-dibenzyl-1,3,5-triazacyclohexan-2-one, and 1-substituted 3,5-dinitro-4-pyridonyl.

## SPECIAL – NH PROTECTIVE GROUPS

5           Examples of special NH protective groups include

### N-Alkyl and N-Aryl Amines

N-methyl, N-allyl, N-[2-(trimethylsilyl)ethoxy]methyl, N-3-acetoxypentyl,  
N-(1-isopropyl-4-nitro-2-oxo-3-pyrrolin-3-yl), quaternary ammonium salts, N-  
10 benzyl, N-di(4-methoxyphenyl)methyl, N-5-dibenzosuberonyl, N-triphenylmethyl,  
N-(4-methoxyphenyl)diphenylmethyl, N-9-phenylfluorenyl, N-2,7-dichloro-9-  
fluorenylmethylene, N-ferrocenylmethyl, and N-2-picolylamine N'-oxide.

### Imine Derivatives

15           N-1,1-dimethylthiomethylene, N-benzylidene, N-*p*-methoxybenzylidene,  
N-diphenylmethylene, N-[(2-pyridyl)mesityl]methylene, and N-(N',N'-  
dimethylaminomethylene).

## 20           PROTECTION FOR THE CARBOXYL GROUP

### Substituted Methyl Esters

Examples of substituted methyl esters include 9-fluorenylmethyl,  
methoxymethyl, methylthiomethyl, tetrahydropyranyl, tetrahydrofuranyl,  
25 methoxyethoxymethyl, 2-(trimethylsilyl)ethoxymethyl, benzyloxymethyl,  
phenacyl, *p*-bromophenacyl,  $\alpha$ -methylphenacyl, *p*-methoxyphenacyl,  
carboxamidomethyl, and N-phthalimidomethyl.

### 2-Substituted Ethyl Esters

30           Examples of 2-substituted ethyl esters include 2,2,2-trichloroethyl,  
2-haloethyl,  $\omega$ -chloroalkyl, 2-(trimethylsilyl)ethyl, 2-methylthioethyl, 1,3-  
dithianyl-2-methyl, 2-(*p*-nitrophenylsulfenyl)ethyl, 2-(*p*-toluenesulfonyl)ethyl,

2-(2'-pyridyl)ethyl, 2-(diphenylphosphino)ethyl, 1-methyl-1-phenylethyl, *t*-butyl, cyclopentyl, cyclohexyl, allyl, 3-buten-1-yl, 4-(trimethylsilyl)-2-buten-1-yl, cinnamyl,  $\alpha$ -methylcinnamyl, phenyl, *p*-(methylmercapto)phenyl and benzyl.

## 5      Substituted Benzyl Esters

Examples of substituted benzyl esters include triphenylmethyl, diphenylmethyl, bis(*o*-nitrophenyl)methyl, 9-anthrylmethyl, 2-(9,10-dioxo)anthrylmethyl, 5-dibenzosuberyl, 1-pyrenylmethyl, 2-(trifluoromethyl)-6-chromylmethyl, 2,4,6-trimethylbenzyl, *p*-bromobenzyl, *o*-nitrobenzyl, *p*-nitrobenzyl, *p*-methoxybenzyl, 2,6-dimethoxybenzyl, 4-(methylsulfinyl)benzyl, 4-sulfobenzyl, piperonyl, 4-picolyl and *p*-P-benzyl.

## 15      Silyl Esters

Examples of silyl esters include trimethylsilyl, triethylsilyl, *t*-butyldimethylsilyl, *i*-propyldimethylsilyl, phenyldimethylsilyl and di-*t*-butylmethylsilyl.

## 20      Activated Esters

Examples of activated esters include thiols.

## 25      Miscellaneous Derivatives

Examples of miscellaneous derivatives include oxazoles, 2-alkyl-1,3-oxazolines, 4-alkyl-5-oxo-1,3-oxazolidines, 5-alkyl-4-oxo-1,3-dioxolanes, ortho esters, phenyl group and pentaaminocobalt(III) complex.

## 30      Stannyl Esters

Examples of stannyl esters include triethylstannyl and tri-*n*-butylstannyl.

## 35      AMIDES AND HYDRAZIDES

### 30      Amides


Examples of amides include N,N-dimethyl, pyrrolidinyl, piperidinyl, 5,6-dihydrophenanthridinyl, o-nitroanilides, N-7-nitroindolyl, N-8-Nitro-1,2,3,4-tetrahydroquinolyl, and *p*-P-benzenesulfonamides.

5      Hydrazides

Examples of hydrazides include N-phenyl and N,N'-diisopropyl hydrazides.



## E. Chemical Examples

Example 1

2-[4-[3-piperidinopropoxy]phenyl]-1-(methanesulfonyl)-1H-indole  
 $K_i = 7 \text{ nM}$

10    Step A        Preparation of 2-iodo-N-(methanesulfonyl)aniline

Methanesulfonyl chloride (4.4 mL, 57 mmol) was added to a 0°C dichloromethane (200 mL) solution containing 2-iodoaniline (5.0 g, 23 mmol) and triethylamine (9.6 mL, 69 mmol). The resulting mixture was stirred for 90 minutes, washed with HCl (0.5 M). The organics were separated then dried over sodium sulfate and concentrated in vacuo. The residue was then treated with potassium hydroxide (2.0 M in 1:1 methanol:water, 75 mL) for 30 min. This material was then diluted with dichloromethane and washed with HCl (1.0 N, 300 mL). The organics were separated then dried over sodium sulfate and concentrated to provide the title compound (5.15 g) as a tan solid.

Step B        Preparation of 4-(methoxyethoxymethyl)benzaldehyde

Sodium hydride (2.4g (60%), 60 mmol) was added to 4-hydroxybenzaldehyde (6.0 g, 50 mmol) in N, N-dimethylformamide (100 mL). The suspension was stirred for 1 hour and then treated with 2-methoxyethoxymethyl chloride (6.8 mL, 60 mmol), and allowed to stir an additional 16 hours. The reaction was then partitioned between water and ether:ethyl acetate (1:1). The organics were then washed with water (4x), dried (potassium carbonate), and concentrated. The crude materials were then purified by silica gel chromatography (hexanes:ethyl acetate) to afford the title compound (9.0 g)

Step C        Preparation of 1-ethynyl-4-(methoxyethoxymethyl)benzene

Dimethyl[(2-diazo-3-oxo)propyl] phosphonate was added in 4 portions to a suspension of potassium carbonate (4.96 g, 36 mmol), the product of Step B (3.78 g, 18 mmol), and methanol (50 mL). The reaction was stirred for 16 hours. and concentrated in vacuo. The residue was taken up in ether, washed  
5 with water (3x), dried (potassium carbonate), and concentrated. The crude product was purified by silica gel chromatography (hexane:ethyl acetate) to provide the title compound (2.3 g).

10 Step D Preparation of 2-(4-(methoxyethoxymethyl)phenyl)-1-(methanesulfonyl)indole

The products of Step A (3.0 g, 10 mmol) and step C (2.2 g, 10 mmol) were combined in N, N-dimethylformamide (20 mL) and triethylamine (5 mL). The solution was then treated with dichlorobis(triphenylphosphine)palladium(II) (0.7  
15 g, 1.0 mmol); copper(I)iodide (380 mg, 2.0 mmol), and stirred at 80 °C for 17 hours. The reaction was then diluted with ether:ethyl acetate (1:1, 200 mL), washed with water (5x), dried (potassium carbonate), and concentrated in vacuo. The crude material was then purified by silica gel chromatography (hexane:ethyl acetate) to afford the title compound (3.36 g).

20 Step E Preparation of 2-(4-hydroxyphenyl)-1-(methanesulfonyl)-indole


A solution of the product of Step D (1.5 g, 4.0 mmol) in methanol (10 mL) was treated with HCl (10 mL, 4 N in dioxane). The reaction was allowed to stir for 2  
25 hr, concentrated, and purified by silica gel chromatography (methanol:dichloromethane), to afford the title compound (0.93 g).

30 Step F Preparation of 2-[4-[3-piperidinopropoxy]phenyl]-1-(methanesulfonyl)indole

A mixture of immobilized triphenylphosphine resin (330 mg, 1.0 meq (Fluka)), and the product of Step E (140 mg, 0.50 mmol) in tetrahydrofuran (6.0 mL) was treated with 3-(piperidin-1-yl)propanol (143 mg, 1.0 mmol) followed by

diethylazidodicarboxylate (0.16 mL, 1.1 mmol). The reaction was shaken for 20 hr. and filtered. The filtrate was concentrated in vacuo and the residue purified by silica gel chromatography (methanol/ethyl acetate) to afford pure title compound (97 mg). MS (ESI)  $m/z$  413 ( $MH^+$ );  $^1H$ -NMR ( $CDCl_3$ )  $\delta$  8.13 (d, 1H), 7.59 (d, 1H), 7.50 (d, 2H), 7.36 (m, 2H), 6.97 (d, 2H), 6.68 (s, 1H), 4.08 (t, 2H), 2.72 (s, 3H), 2.4 (m, 6H), 1.63 (m, 6H), 1.45 (m, 2H).


### Example 2



2-[4-[3-Piperidinopropoxy]phenyl]-1H-indole  
 $K_i = 48$  nM


A solution of the product from Step F, Example 1 (41.4 mg, 0.10 mmol) in methanol (2.0 mL) was treated with potassium hydroxide (1.0 mL, 40% aq). The reaction was stirred at 40° C for 48 hours and concentrated in vacuo. The residue was purified by silica gel chromatography (methanol:dichloromethane) to provide pure title compound (3.7 mg). MS (ESI)  $m/z$  335 ( $MH^+$ );  $^1H$ -NMR ( $CDCl_3$ )  $\delta$  8.19 (bs, 1H), 7.48 (d, 2H), 7.27 (m, 2H), 7.07 (t, 1H), 7.00 (t, 1H), 6.86 (d, 2H), 6.61 (s, 1H), 3.96 (t, 2H), 2.5 (m, 6H), 1.4 (m, 8H).

### Example 3




2-[4-[2-[1-(methyl)-2-pyrrolidinyl]ethoxy]phenyl]-1-(methylsulfonyl)-1H-indole  
 $K_i = 77$  nM

The title compound was obtained (70 mg) by the same general method as Example 1 by substituting 2-ethoxy-1-methylpyrrolidine for 3-(piperdin-1-yl)propanol in step F. MS (ESI)  $m/z$  399 ( $MH^+$ );  $^1H$ -NMR ( $CDCl_3$ )  $\delta$  8.13 (d, 1H), 7.60 (d, 1H), 7.50 (2, 2H), 7.37 (m, 2H), 6.97 (d, 2H), 6.68 (s, 1H), 4.08 (m, 2H), 3.12 (m, 1H), 2.72 (s, 3H), 2.39 (s, 3H), 2.1-1.7 (m, 8H).

Example 4


2-[4-[2-[1-(methyl)-2-pyrrolidinyl]ethoxy]phenyl]-1H-indole  
 $K_i = 100 \text{ nM}$

- The title compound was obtained (14.3 mg) by the same general method as Example 2 by substituting the product of Example 3 for the product of example 1 as the starting material. MS (ESI)  $m/z$  321 ( $MH^+$ );  $^1H$ -NMR ( $CDCl_3$ )  $\delta$  8.15 (bs, 1H), 7.49 (d, 1H), 7.47 (d, 2H), 7.26 (d, 1H), 7.04 (t, 1H), 6.98 (t, 1H), 6.85 (d, 2H), 6.59 (s, 1H), 3.95 (m, 2H), 2.96 (t, 1H), 2.23 (s, 3H), 2.1-1.7 (m, 8H).

Example 5


2-[4-[1-(methyl)-4-piperidinyl]oxyphenyl]-1-(methylsulfonyl)-1H-indole  
 $K_i = 107 \text{ nM}$

- The title compound was obtained (54.8 mg) by the same general method as Example 1 by substituting 4-hydroxy-1-methylpiperidine for 3-(1-piperidinyl) propanol in Step F. MS (ESI)  $m/z$  385 ( $MH^+$ );  $^1H$ -NMR ( $CDCl_3$ )  $\delta$  8.13 (d, 1H), 7.59 (d, 1H), 7.50 (d, 2H), 7.37 (m, 2H), 6.96 (d, 2H), 6.68 (s, 1H), 4.41 (m, 1H), 2.76 (m, 2H), 2.73 (s, 3H), 2.34 (s, 3H), 2.07 (m, 2H), 1.92 (m, 2H), 1.78 (m, 2H).

Example 6


2-[4-[1-(methyl)-4-piperidinyl]oxyphenyl] 1H-indole  
 $K_i = 390 \text{ nM}$

The title compound was obtained (13.8 mg) by the same general method as Example 2 by substituting the product of Example 5 for the product of example 52 as the starting material. MS (ESI)  $m/z$  307 ( $MH^+$ );  $^1H$ -NMR ( $CDCl_3$ )  $\delta$  8.36 (bs, 1H), 7.63 (d, 1H), 7.60 (d, 2H), 7.41 (d, 1H), 7.19 (t, 1H), 7.13 (t, 1H), 7.00 (d, 2H), 6.73 (s, 1H), 4.42 (m, 1H), 2.76 (m, 2H), 2.40 (m, 2H), 2.36 (s, 3H), 2.09 (m, 2H), 1.92 (m, 2H).

Example 7

2-[4-[3-Dimethylaminopropoxy]phenyl]-1-(methylsulfonyl)-1H-indole  
 $K_i = 120$  nM


The title compound was obtained (95 mg) by the same general method as Example 1 by substituting N, N-dimethyl-3-amino-1-propanol for 3-(piperidin-1-yl) propanol in Step F. MS (ESI)  $m/z$  373 ( $MH^+$ );  $^1H$ -NMR ( $CDCl_3$ )  $\delta$  8.13 (d, 1H), 7.59 (d, 1H), 7.50 (d, 2H), 7.36 (m, 2H), 6.97 (d, 2H), 6.68 (s, 1H), 4.09 (t, 2H), 2.73 (s, 3H), 2.50 (t, 2H), 2.29 (s, 6H), 2.01 (m, 2H).

Example 8

2-[4-[3-Dimethylaminopropoxy]phenyl] 1H-indole  
 $K_i = 500$  nM

The title compound was obtained (13.8 mg) by the same general method as Example 2 by substituting the product of Example 7 for the product of Example 1 as the starting material. MS (ESI)  $m/z$  295 ( $MH^+$ );  $^1H$ -NMR ( $CDCl_3$ )  $\delta$  8.10 (bs, 1H), 7.62 (d, 1H), 7.60 (d, 2H), 7.40 (d, 1H), 7.19 (t, 1H), 7.13 (t, 1H), 7.00 (d, 2H), 6.72 (s, 1H), 4.09 (t, 2H), 2.50 (t, 2H), 2.29 (s, 6H), 2.01 (m, 2H).


Example 9



5                                      2-[4-[4-Pyridinyl]methoxyphenyl]-1-(methylsulfonyl)-1H-indole  
 $K_i = 5000 \text{ nM}$

The title compound was obtained (185 mg) by the same general method as  
 Example 1 by substituting 4-hydroxymethylpyridine for 3-(piperdin-1-yl)  
 10    propanol in Step F. MS (ESI)  $m/z$  379 ( $MH^+$ );  $^1H$ -NMR ( $CDCl_3$ )  $\delta$  8.67 (d, 2H),  
 8.13 (d, 1H), 7.60 (d, 1H), 7.54 (d, 2H), 7.41 (d, 2H), 7.38 (m, 2H), 7.03 (d, 2H),  
 6.69 (s, 1H), 4.77 (s, 2H), 2.74 (s, 3H).


Example 10



15                                      2-[4-[2-Diethylaminoethoxy]phenyl]-1-(methylsulfonyl)-1H-indole  
 20                                       $K_i = 369 \text{ nM}$

The title compound was obtained (140 mg) by the same general method as  
 Example 1 by substituting 2-(N, N-diethylamino)ethanol for 3-(piperdin-1-yl)  
 propanol in Step F. MS (ESI)  $m/z$  387 ( $MH^+$ );  $^1H$ -NMR ( $CDCl_3$ )  $\delta$  8.14 (d, 1H),  
 25    7.59 (d, 1H), 7.50 (d, 2H), 7.16 (m, 2H), 6.97 (d, 2H), 6.67 (s, 1H), 4.12 (t, 2H),  
 2.91 (t, 2H), 2.73 (s, 3H), 2.67 (q, 4H), 1.10 (t, 6H).


Example 11



30                                      2-[4-[2-Diethylaminoethoxy]phenyl]-1H-indole  
 35                                       $K_i = 523 \text{ nM}$

The title compound from Example 10 (38.6 mg, 0.10 mmol) was treated with tetrabutyl ammonium fluoride (4.0 mL, of a 0.5 M in tetrahydrofuran) and stirred for 14 hours at 60°C. The resulting solution was concentrated in vacuo, dissolved in dichloromethane, and washed with water. The organics were then concentrated, and the crude product purified by silica gel chromatography (methanol/ dichloromethane) to afford pure title compound (5.9 mg). MS (ESI)  $m/z$  309 ( $MH^+$ );  $^1H$ -NMR ( $CDCl_3$ )  $\delta$  8.21 (bs, 1H), 7.53 (d, 1H), 7.51 (d, 2H), 7.31 (d, 1H), 7.09 (t, 1H), 7.04 (t, 1H), 6.90 (d, 2H), 6.64 (s, 1H), 4.06 (t, 2H), 2.88 (t, 2H), 2.63 (q, 4H), 1.04 (t, 6H).

10

Example 12548/467  
514/414


15

2-[4-[3-(2-Oxo-pyrrolidino)propoxy]phenyl]-1-(methylsulfonyl)-1H-indole

The title compound was obtained (180 mg) by the same general method as Example 1 by substituting 1-(3-hydroxypropyl)-2-pyrrolidone for 3-(piperidin-1-yl) propanol in Step F. MS (ESI)  $m/z$  435 ( $M+Na$ );  $^1H$ -NMR ( $CDCl_3$ )  $\delta$  8.12 (d, 1H), 7.59 (d, 1H), 7.49 (d, 2H), 7.36 (m, 2H), 6.95 (d, 2H), 6.67 (s, 1H), 4.03 (t, 2H), 3.55-3.35 (m, 6H), 2.72 (s, 3H), 2.40 (m, 2H), 2.06 (m, 2H).

20

25


Example 13

30

2-[4-[2-(2-Pyridinyl)ethoxy]phenyl]-1-(methylsulfonyl)-1H-indole

The title compound was obtained (90 mg) by the same general method as Example 1 by substituting 2-(2-hydroxyethyl)pyridine for 3-(piperidin-1-yl)propanol in Step F. MS (ESI)  $m/z$  393 ( $MH^+$ );  $^1H$ -NMR ( $CDCl_3$ )  $\delta$  8.59 (d, 1H), 8.12 (d, 1H), 7.67 (d, 1H), 7.59 (d, 1H), 7.48 (d, 2H), 7.35 (m, 2H), 7.20 (m, 2H), 6.97 (d, 2H), 6.66 (s, 1H), 4.43 (t, 2H), 3.32 (t, 2H), 2.72 (s, 3H).


35

Example 14

546/277.4  
514/339

2-[4-[2-(2-Pyridinyl)ethoxy]phenyl]-1H-indole

The title compound was obtained (14.5 mg) by the same general method as Example 11 by substituting the product of Example 13 for the product of Example 10 as the starting material. MS (ESI)  $m/z$  315 ( $MH^+$ );  $^1H$ -NMR ( $CDCl_3$ )  $\delta$  8.52 (d, 1H), 8.45 (bs, 1H), 7.58 (t, 1H), 7.53 (d, 1H), 7.50 (d, 2H), 7.29 (d, 1H), 7.22 (d, 1H), 7.12 (t, 1H), 7.08 (t, 1H), 7.01 (t, 1H), 6.88 (d, 2H), 6.63 (s, 1H), 4.32 (t, 2H), 3.21 (t, 2H).


Example 15

546/201  
514/323

2-[3-[3-Piperidinopropoxy]phenyl]-1-(methylsulfonyl)-1H-indole  
 $K_i = 33$  nM

The title compound was obtained (84 mg) by the same general method as Example 1 by substituting 3-hydroxybenzaldehyde for 4-hydroxybenzaldehyde in Step B. MS (ESI)  $m/z$  413 ( $MH^+$ );  $^1H$ -NMR ( $CDCl_3$ )  $\delta$  8.14 (d, 1H), 7.62 (d, 1H), 7.37 (m, 3H), 7.16 (d, 1H), 7.12 (s, 1H), 6.98 (d, 1H), 6.74 (s, 1H), 4.08 (t, 2H), 2.77 (s, 3H), 2.62 (t, 2H), 2.55 (m, 2H), 2.10 (m, 2H), 1.69 (m, 4H), 1.50 (m, 1H).



Example 16

2-(4-(3-(4-Methylpiperazino)propoxy)phenyl)indole

 $K_i = 2000 \text{ nM}$ 

## Step A Preparation of 4'-(3-chloropropoxy)acetophenone

A solution of 4'-hydroxyacetophenone (20 mmol, 2.72 g), 3-bromopropionyl chloride (21 mmol, 2.07 mL) and potassium carbonate (4.14g, 30.0 mmol) in acetone (50 mL) was heated at reflux for overnight. The salt was filtered off. The solvent was evaporated. After drying in vacuo, the title compound (4.24 g) was collected.

## Step B Preparation of 2-(4-(3-chloropropoxy)phenyl)indole


A mixture 4'-(3-chloropropoxy)acetophenone (10 mmol, 2.12 g) and phenylhydrazine (10 mmol, 1.08 g) was heated at 100 °C for 40 min. Then PPA (~5 g) was added and temperature was raised to 130 °C and kept for 10 min. The mixture was cooled down. Water (50 mL) was added. After 2 h, greenish solid was formed and collected via filtration. The solid then was washed by a small amount of methanol (5 mL). After drying in vacuo, the title compound (1.5 g) was obtained.

## Step C 2-(4-(3-(4-Methylpiperazino)propoxy)phenyl)indole

The mixture of 2-(4-(3-chloropropoxy)phenyl)indole (1 mmol, 285 mg) and 4-methylpiperazine (2 mL) was heated at 80 °C for 5 h. After concentration, the residue was purified by column chromatography (MeOH/CH<sub>2</sub>Cl<sub>2</sub>) afforded the title compound (285 mg): <sup>1</sup>H NMR (CDCl<sub>3</sub>, 400MHz) δ 8.26 (s, 1H), 7.53 (m, 3H), 7.32 (bd, 1H, J = 8.1 Hz), 7.1 (td, 1H, J = 1.1, 7.0 Hz), 7.05 (td, 1H, J =

1.1, 7.0 Hz), 6.90 (m, 2H), 6.64 (m, 1H), 3.99 (t, 2H,  $J = 6.3$  Hz), 2.48 (m, 10H,  $J = 6.4$  Hz), 2.25 (s, 3H), 1.93 (quintet, 2H,  $J = 6.3$  Hz); EIMS  $m/z$  350 ( $M + H^+$ ).

5

Example 17

544/373  
514/254.09

1-(Methylsulfonyl)-2-(4-(3-(4-methylpiperazino)propoxy)phenyl)indole  
 $K_i = 3000$  nM

10

**Step A** Preparation of 4-(3-chloropropoxy)benzaldehyde

A solution of 4-hydroxybenzaldehyde (100 mmol, 12.2 g), 3-bromopropionyl chloride (101 mmol, 20 mL) and potassium carbonate (20.7 g, 150 mmol) in acetone (250 mL) was heated at reflux for overnight. The salt was filtered off. The solvent was evaporated. Reduced pressure distillation provided the title compound (15 g).

15

**Step B** Preparation of 1-ethynyl-4-(3-chloropropoxy)benzene

20

To LDA (2M in THF, 15 mL) in THF (100 mL) at  $-78^\circ\text{C}$ ,  $\text{TMSCHN}_2$  (2M in hexanes, 15 mL) was added dropwisely. Ten minutes later, 4-(3-chloropropoxy)benzaldehyde (0.025 mol, 4.97 g) in THF (60 mL) was added. After 1 h stirring at  $-78^\circ\text{C}$ , the mixture was warmed up and refluxed for 3h. Water (250 mL) was added and extracted by EtOAc (2 x 200 mL). After being dried over  $\text{Na}_2\text{SO}_4$  and concentration, the title compound (4.8 g) was obtained.

25

**Step C** Preparation of 1-ethynyl-4-(3-(4-methylpiperazino)propoxy)benzene

30

The mixture of 1-ethynyl-4-(3-chloropropoxy)benzene (2 mmol, 388 mg) and 4-methylpiperazine (2 mL) was heated at 80 °C for 5h. After concentration, the residue was purified by column chromatography (MeOH/CH<sub>2</sub>Cl<sub>2</sub>) afforded the title compound (400 mg).

5

#### Step D Preparation of 2-iodo-N-(methanesulfonyl)aniline

To the mixture of the 2-iodo-4-chloroaniline (5.0 g, 20 mmol) and triethylamine (8.3 mL, 60 mmol) in methylene chloride (200 mL), the solution of methanesulfonyl chloride (3.4 mL, 44 mmol) was added slowly. The solution then was stirred at room temperature for 2 h. After being washed by HCl solution (0.5 N, 2 x 100 mL), NaOH solution (0.5 N, 2 x 100 mL), brine (100 mL), the organic layer was dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated providing the title compound (6.6 g).

15

#### Step E Preparation of 1-(methylsulfonyl)-2-(4-(3-(4-methylpiperazino)propoxy)phenyl)indole

The mixture of 1-ethynyl-4-(3-(4-methylpiperazino)propoxy)benzene (230 mg, 0.89 mmol), 2-iodo-N-(methanesulfonyl)aniline (0.89 mmol, 296 mg), CuI (17 mg, 0.089 mmol), Pd(PPh<sub>3</sub>)<sub>2</sub>Cl<sub>2</sub> (32 mg, 0.045 mmol) and triethylamine (0.5 mL) in DMF (5 mL) was stirred at 80 °C for overnight. After concentration, water (20 mL) was added and extracted by methylene chloride (3 x 20 mL). The organics was concentrated and purified by column chromatography afforded the title compound (260 mg): <sup>1</sup>H NMR (CDCl<sub>3</sub>, 400MHz) δ 8.05 (d, 1H, J = 8.8 Hz), 7.55 (d, 1H, J = 2.1 Hz), 7.50 (td, 2H, J = 8.8, 2.1 Hz), 7.32 (d, 1H, J = 2.1 Hz), 7.30 (d, 1H, J = 2.1 Hz), 6.96 (td, 2H J = 2.1, 8.8 Hz), 6.60 (d, 1H, J = 0.5 Hz), 4.08 (t, 2H, J = 6.3 Hz), 2.74 (s, H), 2.55 (m, 10H), 2.33 (s, 3H), 2.02 (quintet, 2H, J = 6.3 Hz); EIMS m/z 462 (M + H<sup>+</sup>).

30

## F. Biological Examples

Biological Example 1

## 1(A) Transfection of cells with human histamine receptor

- 5           A 10 cm tissue culture dish with a confluent monolayer of SK-N-MC cells was split two days prior to transfection. Using sterile technique the media was removed and the cells were detached from the dish by the addition of trypsin. One fifth of the cells were then placed onto a new 10 cm dish. Cells were grown in a 37°C incubator with 5% CO<sub>2</sub> in Minimal Essential Media Eagle with
- 10   10% Fetal Bovine Serum. After two days cells were approximately 80% confluent. These were removed from the dish with trypsin and pelleted in a clinical centrifuge. The pellet was then re-suspended in 400 µL complete media and transferred to an electroporation cuvette with a 0.4cm gap between the electrodes (Bio-Rad #165-2088). One µg supercoiled H<sub>3</sub> receptor cDNA
- 15   was added to the cells and mixed. The voltage for the electroporation was set at 0.25 kV, the capacitance is set at 960 µF.

- After electroporation the cells were diluted into 10 mL complete media and plated onto four 10cm dishes. Due to the variability in the efficiency of electroporation, four different concentrations of cells were plated. The ratios
- 20   used were: 1:20, 1:10, and 1:5, with the remainder of the cells being added to the fourth dish. The cells were allowed to recover for 24 hours before adding the selection media (complete media with 600 µg/ml G418). After 10 days dishes were analyzed for surviving colonies of cells. Dishes with well-isolated colonies were used. Cells from individual colonies were isolated and tested.
- 25   SK-N-MC cells were used because they give efficient coupling for inhibition of adenylate cyclase. The clones that gave the most robust inhibition of adenylate cyclase in response to histamine were used for further study.

## 1(B) [3H]-N-methylhistamine binding

- 30           Cell pellets from histamine H<sub>3</sub> receptor-expressing SK-N-MC cells were homogenized in 20 mM TrisHCl/0.5mM EDTA. Supernatants from a 800 g spin were collected, re-centrifuged at 30,000 g for 30 minutes. Pellets were rehomogenized in 50 mM Tris/5mM EDTA (pH 7.4). Membranes were

- incubated with 0.8 nM [<sup>3</sup>H]-N-methylhistamine plus/minus test compounds for 45 minutes at 25°C and harvested by rapid filtration over GF/C glass fiber filters (pretreated with 0.3% polyethylenimine) followed by four washes with ice cold buffer. Filters were dried, added to 4 mL scintillation cocktail and then
- 5 counted on a liquid scintillation counter. Non-specific binding was defined with 10 µM histamine. PK<sub>i</sub> values were calculated based on a K<sub>D</sub> of 800 pM and a ligand concentration ([L]) of 800 pM according to the formula:

$$K_i = (IC_{50}) / (1 + ([L] / K_D))$$

10

11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000

# G. Other Embodiments

The features and advantages of the invention are apparent to one of ordinary skill in the art. Based on this disclosure, including the summary, detailed description, background, examples, and claims, one of ordinary skill in the art will be able to make modifications and adaptations to various conditions and usages. These other embodiments are also within the scope of the invention.

10

15

What is claimed is: